


Longitudinal Spin Transfer to Hyperons in Polarized p+p Collisions at $\sqrt{s}=200$ GeV

Ernst Sichtermann (LBNL), *for the*  *STAR* Collaboration

Nucleon Spin Puzzle

The surprising *smallness* of the spin dependent part of the inclusive DIS cross section renewed the interest in nucleon spin structure,



EMC (1988): Quark and anti-quark spins combined contribute little to the proton spin, Strange (anti-)quarks are negatively polarized.


Among the many open questions, what is the role of *strange (anti-)quark spins*, is there a *hyperon spin puzzle*?

What insight(s) can hyperon polarization measurements at RHIC give?


Nucleon Spin Puzzle

Semi-inclusive DIS data with identified Kaons in the final state add precision, and pose yet more questions:

COMPASS, Phys.Lett.B693 (2010) 227



COMPASS, Phys.Lett.B680 (2009) 217



$$R_{SF} = \frac{\int D_s^{K^+}(z)dz}{\int D_u^{K^+}(z)dz}$$



Clear call for complementary measurements,

Notoriously hard at RHIC; charm-associated W production (Sudoh, 2005 RHIC-II w.s.),
Try hyperon spin-transfer.

RHIC - Polarized Proton Collider to Study Spin in QCD

Opportunities to study many facets:

$\sqrt{s} = 200 - 500 \text{ GeV}$




with good systematic controls, e.g.:




This talk: $\sqrt{s} = 200 \text{ GeV}$, $\sim 3 \text{ pb}^{-1}$, $P_b \sim 50\%$ (longitudinal), collected in Y2005
 $\sim 22 \text{ pb}^{-1}$, $P_b \sim 57\%$ (longitudinal), collected in Y2009


STAR - Solenoid Tracker At RHIC



Time Projection Chamber enables PID,




and topological reconstruction,




for $|\eta| \leq \sim 1.3$

Differential Cross Section



Factorized framework,




$$f, \Delta f \otimes \hat{\sigma}, \Delta \hat{\sigma} \otimes D, \Delta D$$

enables perturbative description.


Agreement of STAR data and theory, for a *suitable* choice of D , is a necessary condition for interpretation.

Note: The AKK 2008 update again undershoots the STAR data,
 Opportunities exist also to extend the data to higher p_T (eventually).

Spin-dependent Fragmentation



Factorized framework,



$$f, \Delta f \otimes \hat{\sigma}, \Delta \hat{\sigma} \otimes D, \Delta D$$

enables perturbative description.

Polarized fragmentation is sizable, especially for large fragmentation momentum-fractions z ,

Note: data remain scarce,
 ΔD is thus often *modeled*.

D_{LL} - Longitudinal Spin Transfer

At RHIC,

$$D_{LL}^{\Lambda} \equiv \frac{\sigma_{p^+ p \rightarrow \Lambda^+ X} - \sigma_{p^+ p \rightarrow \Lambda^- X}}{\sigma_{p^+ p \rightarrow \Lambda^+ X} + \sigma_{p^+ p \rightarrow \Lambda^- X}} = P_{\Lambda}^+$$

that is, the longitudinal polarization of the Λ for a specific beam-helicity configuration.

This polarization can be determined in the usual way,

$$\frac{dN}{d\Omega} \propto A(\cos \theta^*) (1 + \alpha P_{\Lambda} \cos \theta^*)$$


from the angular distribution of the $p + \pi$ decay mode with B.R. $\sim 64\%$.

Here,


A is the detector acceptance (which can be canceled in a ratio analysis),
 θ^* is the angle defined by the Λ momentum and the p direction in the Λ rest frame,
 $\alpha = 0.642 \pm 0.013$ is the decay parameter.

D_{LL} - Longitudinal Spin Transfer

Expectations at LO show sensitivity of D_{LL} for the $\bar{\Lambda}$ to the \bar{s} helicity distribution, $\Delta\bar{s}$,



GRSV00 - M. Glück et al Phys.Rev.D63 (2001) 094005




more so than to the fragmentation in this model.

The Λ D_{LL} is less sensitive to Δs , partly due to larger u and d quark fragmentation contributions.

Promising measurement: neither the role of (anti-)strange quarks nor polarized fragmentation is well known/understood - effects are potentially large enough to be observed.

For the Spin-Aficionados - Measure D_{LL} or A_{LL} ?

The same expectations versus p_T as D_{LL} and A_{LL} :



$$D_{LL} \equiv \frac{\sigma_{p^+ p \rightarrow \Lambda^+ X} - \sigma_{p^+ p \rightarrow \Lambda^- X}}{\sigma_{p^+ p \rightarrow \Lambda^+ X} + \sigma_{p^+ p \rightarrow \Lambda^- X}}$$



$$A_{LL} \equiv \frac{\sigma_{p^+ p^+ \rightarrow \Lambda X} - \sigma_{p^+ p^- \rightarrow \Lambda X}}{\sigma_{p^+ p^+ \rightarrow \Lambda X} + \sigma_{p^+ p^- \rightarrow \Lambda X}}$$

- + D_{LL} expected sensitivity is ~ 4 larger,
- D_{LL} analysis requires more selections than for A_{LL} , i.e. lose some statistics,
- + D_{LL} is a single beam-spin measurement, analyzing power of the $p+\pi$ decay mode is relatively large.

Net advantage owing to the (anti-) Λ spin being carried mostly by the (anti-) s quark spin.

STAR Initial Data - 2005

$\sim 3 \cdot 10^6$ *minimum bias* events (beam-collision triggered, band-width limited),



$\sim 30 \cdot 10^3$ Λ candidates,

$\sim 25 \cdot 10^3$ $\bar{\Lambda}$



$\langle p_T \rangle \approx 1.3$ GeV/c

$\langle |x_F| \rangle \approx 0.008$

Take away: analyze data triggered on hard-processes...

STAR Triggered Data - 2005

STAR was triggered on energy deposits in jet-patches of the Barrel E.M. Calorimeter,




Although this is not a “Hyperon Trigger”, it did record a (biased) sample of Λ and $\bar{\Lambda}$ candidates with considerably higher p_T ; focus on $\bar{\Lambda}$ here.

Analysis Characteristics

Uses the $\Lambda \rightarrow p + \pi$ weak decay mode,

$$\frac{dN}{d\Omega} \propto A(\cos \theta^*) (1 + \alpha P_\Lambda \cos \theta^*)$$

Restrict $\cos \theta^*$ to eliminate K_S^0 background caused by misidentified π , (refined in later analyses).




Use beam spin configurations and symmetries to (largely) cancel $A(\cos \theta^*)$ and extract,

$$D_{LL}^\Lambda = \frac{1}{\alpha \cdot P_b \cdot \langle \cos \theta^* \rangle} \cdot \frac{N_\Lambda^+ - N_\Lambda^-}{N_\Lambda^+ + N_\Lambda^-}$$

in small $\cos \theta^*$ intervals. Here, $N_\Lambda^+ = N_\Lambda^{++} \cdot \frac{\mathcal{L}^{--}}{\mathcal{L}^{++}} + N_\Lambda^{+-} \cdot \frac{\mathcal{L}^{--}}{\mathcal{L}^{+-}}$ and $N_\Lambda^- = N_\Lambda^{-+} \cdot \frac{\mathcal{L}^{--}}{\mathcal{L}^{-+}} + N_\Lambda^{--}$

The luminosity ratios are measured at STAR and beam polarization in RHIC.

STAR Initial Results - 2005



D_{LL} proof-of-concept from RHIC,

Systematics under control,


Statistics limited,

0.08 at $p_T \sim 4$ GeV/c

$\langle p_T \rangle \approx 1.3$ GeV/c, $\langle |x_{Fl}| \rangle \approx 0.008$

Take away: need better precision and higher p_T

STAR - 2009



Full-coverage Barrel EMCal,

Trigger improvement,


DAQ-1000,

RHIC luminosity and polarization, even though
the run was cut short and FoM remained
a factor below our initial projections,

Good reasons for continued 200 GeV!

R. Cendejas (UCLA/LBNL) PhD Thesis 2012,
J. Deng (Shandong U.)



STAR - 2009



$$\Delta R = \sqrt{\Delta\phi^2 + \Delta\eta^2}, \Delta R < 0.7$$


$$\Delta\phi = \phi_{\Lambda} - \phi_{jet}$$

$$\Delta\eta = \eta_{\Lambda} - \eta_{jet}$$



Decision to focus, at least initially, on Hyperons that are part of the near-side (trigger) jet. 16

STAR - 2009






Systematic uncertainties vary from 0.01 to 0.03 for each point which include:

- 4.7% Beam polarization
- 2.0% Decay parameter
- 1.9% Residual trans. pol.
- 5×10^{-3} Relative luminosity
- $< 6 \times 10^{-3}$ Residual background.
- ≤ 0.03 Trigger bias, increases with p_T .
- ≤ 0.01 Pile-up, decreases with p_T .


J. Deng for the collaboration, SPIN 2012
R. Cendejas for the collaboration, DNP 2012

Some Cross-Checks



Measurements with the expected null-results.

Compared to Published Results




Systematic uncertainties vary from 0.01 to 0.03 for each point which include:

- 4.7% Beam polarization
- 2.0% Decay parameter
- 1.9% Residual trans. pol.
- 5×10^{-3} Relative luminosity
- $< 6 \times 10^{-3}$ Residual background.
- ≤ 0.03 Trigger bias, increases with p_T .
- ≤ 0.01 Pile-up, decreases with p_T .

J. Deng for the collaboration, SPIN 2012
 R. Cendejas for the collaboration, DNP 2012

D_{LL} out to $p_T \sim 5.9$ GeV with $\sim 4\%$ precision (2009), c.f. $\sim 8\%$ at 3.7 GeV published (2005).

Compared to Expectations - I





D. de Florian, M. Stratmann,
and W. Vogelsang, PRL. 81.
(updated calculation to low p_T)

scen. 1: SU(6) picture.
scen. 2: DIS picture.
scen. 3: equal contribution.

J. Deng for the collaboration, SPIN 2012
R. Cendejas for the collaboration, DNP 2012


Compared to Expectations - II



Data do not currently discriminate “model”-expectations,
precision may re-interest our theory friends, “models” → fits
analysis of away-side sample in progress.

Looking Ahead

- H.Z. Huang for the collaboration, QM2012




- Forward instrumentation optimized for **p+A** and **transverse spin** physics
 - Charged-particle tracking
 - e/h and γ/π^0 discrimination
 - Possibly Baryon/meson separation

- TPC inner sector upgrade will extend acceptance to larger rapidity where D_{LL} is expected to be larger.

- Forward Calorimeter upgrade

W. Zhou et al, Phys.Rev.D 81 (2010) 057501

$\sqrt{s} = 500$ GeV, $p_T > 3$ GeV/c, $2.5 < \eta < 3.5$



should enable also other physics with (forward) Hyperons.

- STAR Decadal Plan discussed in Jamie's talk tomorrow,

- Stay Tuned, Thanks!